

ALTERNATIVE CATALYST MANUFACTURING METHODS

Completed February 2011
(for the 2010 membership year)

*A technical investigation commissioned by the members of the **Catalytic Advances Program***

CONTENTS

| | |
|---|-----------|
| EXECUTIVE SUMMARY | IX |
| 1. INTRODUCTION | 1 |
| 2. NEW MATERIALS AND SYNTHESIS TECHNOLOGIES | 5 |
| 2.1 OXIDES, CARBIDES, NITRIDES, AND SULFIDES | 5 |
| 2.2 CARBON NANOTUBES AND THEIR MODIFICATION | 13 |
| 2.2.1 ODH of hydrocarbons | 14 |
| 2.2.2 ORR in electrocatalysis | 15 |
| 2.2.3 Nanoparticle confinement in CNTs and catalysis | 17 |
| 2.3 ZEOLITES AND ZEOTYPES | 19 |
| 2.3.1 Zeolites and other inorganic molecular sieves | 19 |
| 2.3.2 Zeolitic imidazolate frameworks | 21 |
| 2.4 METAL-ORGANIC FRAMEWORKS | 24 |
| 2.5 ORDERED MESOPOROUS MATERIALS | 28 |
| 2.6 NANOSIZED METALS AND ALLOYS | 30 |
| 2.7 PEROVSKITES AND OTHER “INTELLIGENT” CATALYTIC MATERIALS | 34 |
| 2.8 STATUS OF COMMERCIAL PRACTICALITY AND HURDLES | 38 |
| 2.9 FUTURE NEEDS | 40 |
| 2.10 REFERENCES | 41 |
| 3. NEW METHODS OF CATALYST FORMULATION | 51 |
| 3.1 METALS AND ACTIVE COMPONENT INCORPORATION | 51 |
| 3.1.1 Electroless deposition | 51 |

| | | |
|-------|---|-----|
| 3.1.2 | Chemical vapor deposition..... | 53 |
| 3.1.3 | Atomic layer deposition | 54 |
| 3.2 | SPECIFIC CONTROL ON THE ATOMIC SCALE..... | 56 |
| 3.2.1 | Particle size and distribution control | 56 |
| 3.2.2 | Crystallite size and shape control..... | 58 |
| 3.3 | NEW FUNCTIONALIZATION METHODOLOGIES..... | 62 |
| 3.3.1 | Control of surface acidity and accessibility of acid sites | 62 |
| 3.3.2 | Mixed metal oxides | 64 |
| 3.3.3 | Addition of other inorganic modifiers..... | 65 |
| 3.3.4 | Immobilization of homogeneous catalysts..... | 65 |
| 3.3.5 | Organic modifiers..... | 67 |
| 3.4 | NEW METHODS OF ACTIVATION..... | 69 |
| 3.5 | CATALYST RECOVERY AND REUSE | 70 |
| 3.5.1 | Strategies for maintaining metal distributions of near-surface alloys..... | 70 |
| 3.5.2 | Strategies for maintaining nanoparticle shape and size..... | 71 |
| 3.5.3 | Strategies for maintaining coverage by functionalizing compounds | 73 |
| 3.6 | ASSESSMENT OF COMMERCIAL REALITY, GAPS AND NEEDS | 73 |
| 3.7 | REFERENCES..... | 75 |
| 4. | NEW FORMS OF CATALYSTS | 83 |
| 4.1 | PHASE TRANSFER CATALYSTS..... | 83 |
| 4.2 | MICELLAR CATALYSTS | 88 |
| 4.2.1 | Basic structure, physical and chemical properties of micelles..... | 88 |
| 4.2.2 | Micellar catalysis – catalysis within micelles | 89 |
| 4.2.3 | Micellar catalysts – utilizing micelles to synthesize catalysts and catalyst precursors | 91 |
| 4.3 | IONIC LIQUIDS..... | 94 |
| 4.3.1 | Synthesis and catalysis of supported ionic liquids (heterogenized homogeneous catalysts)..... | 95 |
| 4.3.2 | Catalysis of metal nanoparticles in ionic liquids..... | 98 |
| 4.3.3 | Homogeneous (organometallic) catalysis in ionic liquids | 102 |
| 4.3.4 | Enzymatic (bio) catalysis | 103 |
| 4.3.5 | Brønsted and Lewis acid catalysis in ionic liquids..... | 104 |

| | |
|---|------------|
| 4.4 OTHER NOVEL CATALYSTS | 106 |
| 4.4.1 Metal foams | 106 |
| 4.4.2 Metal organic framework materials | 109 |
| 4.4.3 Multi-component catalysts | 112 |
| 4.5 CATALYST RECOVERY AND REUSE | 117 |
| 4.6 FACTORS IMPACTING LARGE SCALE UTILIZATION | 118 |
| 4.7 STATUS OF COMMERCIAL PRACTICALITY, HURDLES, FUTURE NEEDS | 119 |
| 4.8 REFERENCES | 120 |
| 5. INDEX | 129 |

FIGURES

| | | |
|------------|--|----|
| Figure 2.1 | CdSe nanorods (from Thoma et al., 2005) | 12 |
| Figure 2.2 | PdCoSx nanoacorns (from Teranishi et al., 2004). | 13 |
| Figure 2.3 | (left) Layered structure of g-C ₃ N ₄ ; (right) graphite-like stacking in g-C ₃ N ₄ . The power spectra of areas 1 and 2 show 0.327 nm <i>d</i> -spacing and streaks due to stacking faults (from Thomas et al., 2008)..... | 14 |
| Figure 2.4 | 3D TEM of CNT-NP System: Blue Pt–Ru NPs outside, Red Pt–Ru NPs inside, and Yellow Au NPs Deposited on the CNT Surface for Image Reconstruction (from Serp and Castillejos, 2010)..... | 18 |
| Figure 2.5 | The bridging angles in ZIFs (1) and zeolites (2) (from Park et al., 2006). | 21 |
| Figure 2.6 | (left) Bright-field TEM Image of a Au@ZIF-8 crystal; (right) 3D Rendered Volume Showing the Au NP Distribution and the Edges of the Imaged Crystal (from Esken et al., 2010)..... | 23 |
| Figure 2.7 | Flow Diagram of Industrial MOF Synthesis via Solvothermal (left, Mg-MOF) and Electrochemical (right, Cu-EMOF) Routes (from Czaja et al., 2009) | 25 |
| Figure 2.8 | Self-assembly Pathway of Ordered Mesoporous Metal Oxides (from Carreon and Guliants, 2009a,b)..... | 28 |
| Figure 2.9 | TEM Images of SBA-16 Templated Co ₃ O ₄ (inset: SAED pattern); Image in (d) is Viewed Down the [-110] Axis and Shows (11-1) and (111) Planes. The arrow points to the bridge connecting two Co ₃ O ₄ nanospheres (from Yue and Zhou, 2007b)..... | 30 |

| | | |
|-------------|--|----|
| Figure 2.10 | TEM images of anisotropic Pd nanoparticles: (A) nanorods and polyhedra; (B) a typical nanorod with a five-fold symmetry; (C) a polyhedron formed from six tetrahedral subunits; (D) a tetrahedral particle with rounded edges; (E) nanocube with rounded edges (from Berhault et al., 2007)..... | 32 |
| Figure 2.11 | 3D analysis of a bipyramidal Pd NP: (a) surface rendering of the tomogram; (b) stretched superposition of slices extracted every 6 nm from the tomogram (from Benlekbir et al., 2009)..... | 33 |
| Figure 2.12 | Proposed buta-1,3-diene hydrogenation on Pd NPs. The dotted crosses show unfavorable routes for Pd NPs vs. the isotropic catalyst, whereas solid crosses indicate prohibited routes (from Berhault et al., 2007)..... | 34 |
| Figure 2.13 | Unit Cell of Cubic ABO_3 Perovskite..... | 34 |
| Figure 2.14 | TEM Observation of Pt on $CaTiPtO_3$ after (d) oxidation, (e) reduction, (f) re-oxidation (from Taniguchi et al., 2007)..... | 36 |
| Figure 3.1 | Scheme for electroless deposition illustrating (a) deposition on the support, (b) catalytic deposition through reaction with a reducing agent on the host metal, and (c) autocatalytic deposition through reaction with a reducing agent on already deposited metal. For definiteness, Ag is used as the metal to be deposited by ED, with Pt being the host metal | 52 |
| Figure 3.2 | Schematic for atomic layer deposition of Al_2O_3 from the self-limiting reactions of $Al(CH_3)_3$ and H_2O | 55 |
| Figure 3.3 | Electron microscope images and models of different shapes of Ag metal particles: a, g) right bipyramids; b, h) pentagonal nanowires; c, i) cubes; d, j) truncated right bipyramids; e, k) quasi-spherical; f, l) truncated cubes. Reproduced with permission from (Linic, et al., 2010)..... | 59 |
| Figure 3.4 | Summary of use of polymer networks for preparation of various novel forms of zeolites. Reproduced with permission from (Yao, et al., 2010)..... | 62 |
| Figure 3.5 | Novel approaches for preparation of heterogeneous catalysts from organometallic complexes. Reproduced with permission from (Tada, et al., 2006)..... | 67 |
| Figure 3.6 | Aggregate-free and well-dispersed Au nanoparticles intercalated into the walls of mesoporous silica, from the reference cited below. Reproduced with permission from (Chen, et al., 2009)..... | 72 |
| Figure 4.1 | The mechanism of liquid/liquid phase transfer catalysis with tetraalkylonium salts as phase-transfer catalysts for the production of cyanooctane from chlorooctane. (Ooi, et al., 2007)..... | 84 |

| | | |
|------------|---|----|
| Figure 4.2 | Different types of quaternary ammonium salts for asymmetric alkylation of amino acids including a) binaphthyl amino compounds b) cinchona alkaloid c) tartrate and d) unclassified amino derivatives (Hashimoto, et al., 2007).....85 | 85 |
| Figure 4.3 | a) Maruoka catalyst and simplified Maruoka catalyst for large scale production of synthetic amino acids. b) reaction scheme using the simplified Maruoka catalyst. Personal communication with Professor Keiji Maruoka (10 January 2011).....86 | 86 |
| Figure 4.4 | Schematic of a typical micelle in aqueous solution. In a micelle, the hydrophilic “head” region (purple spheres) of an amphiphile (surfactant) is exposed and in contact with the surrounding solvent (for example, water) and the hydrophobic portion of the amphiphile is sequestered to the interior of the micelle. In a reverse micelle, the head region is sequestered to the interior of the micelle and the tails extend out into solution. Both types of micelles are approximately spherical in shape. From http://www.uic.edu/classes/bios/bios100/lecturesf04am/lect02.htm (accessed on 21 January 2011).89 | 89 |
| Figure 4.5 | Rhodium nanoparticles synthesized by the reduction of Na_3RhCl_6 in butylammonium laurate reverse micelles using sodium borohydride. The molar ratio of surfactant: Na_3RhCl_6 was 30:1 (top), 6:1 (middle) and 4.8: 1 (bottom). The corresponding particle size for each synthesis is shown the histograms. The as-synthesized particles scale with the average micelle size determined by dynamic light scattering (Hoefelmeyer, et al., 2007).92 | 92 |
| Figure 4.6 | (A-C) TEM micrograph images of Pt nanoparticles synthesized by the reverse micelle method and subsequently supported on P25 TiO_2 . (A) Sample #1 before reaction; (B) sample #1 after reaction and (C) sample #3 before reaction. All catalysts were calcined at 773 K. (D) Activity of micelle-derived Pt catalyst after supporting on anatase TiO_2 for the steam reforming of methanol. The catalyst display increasing activity with temperature, and the catalyst with the smallest particle size (Sample #1, $d = 8.6$ nm) is the most active. Samples #3 and #5 have similar activity for steam reforming even though the particle size differs (11.4 nm (sample #3) versus 20.8 nm (sample #5)). (Croy, et al., 2007)94 | 94 |
| Figure 4.7 | Examples of ionic liquids. There are more than one million binary combinations of ionic liquids. Many of their physical properties which are influential to catalysis can be tuned by changing the composition of the organic cation and of the anion. Common ionic liquid cations include (a) imidazolium; (b) thiazolium; (c) pyrazolium; (d) pyridinium; (e) pyrrolinium and (f) oxazolium. Anions can either be inorganic in nature, and include BF_4 , PF_6 , AsF_6 , SbF_6 , SnCl_3 or organic in nature, such as CH_3SO_3 , CF_3CO_2 , $(\text{CF}_3\text{SO}_2)_3\text{C}$, $\text{CF}_3(\text{CF}_2)_2\text{CO}_2$ and $\text{CF}_3(\text{CF}_2)_3\text{SO}_3$. Neither the list of cations or anions should be considered exhaustive.95 | 95 |

| | | |
|-------------|---|-----|
| Figure 4.8 | Schematic of Supported Ionic Liquid Catalyst (SILCs). The Rh pre-catalyst [Rh(NBD)(PPh ₃) ₂]PF ₆ (NBD = norbornadiene, PPh ₃ = triphenylphosphine) dissolved in [bmim][PF ₆], acetone and mixed with silica. Upon removal of the volatile solvent, the SILC catalyst contained an ionic liquid loading of 25%. This catalyst was active for hydrogenation of organic compounds (Mehnert <i>et al.</i> , 2002b). | 96 |
| Figure 4.9 | Preparation of a surface-anchored ionic liquid silica. A highly-selective hydroformylation catalyst was prepared by contacting Rh(CO) ₂ acetylacetonate with the appropriate phosphine-based ligand and appropriate ionic liquid ([bmim][BF ₄] or [bmim][PF ₆]) in acetonitrile and then contacted with the ionic-liquid modified silica. After removal of the volatile solvent, the SILC catalyst contained 25 % by weight ionic liquid (Mehnert, et al., 2002a). | 96 |
| Figure 4.10 | (A) Schematic of immobilized Pd nanoparticles on the surface of SBA-15 mesoporous silica with a layer of ionic liquid on the surface. The ionic liquid – 1,1,3,3-tetramethylguanidinium lactate -- depicted in the figure as + and – signs covers the catalyst as a single monolayer enabling hydrogenation reactions to be conducted under neat, solvent-free conditions. The propose coordination of the Pd nanoparticle to the cation is depicted in (A). (B) TEM micrograph of Pd nanoparticles synthesized by colloidal routes. The particles had a diameter ranging from 1-2 nm. (C) TEM micrograph of Pd nanoparticles loaded into mesoporous SBA-15 silica (Huang, et al., 2004). | 98 |
| Figure 4.11 | (A) C-H borylation of arenes by pinacol borane utilizing ionic acid stabilized Ir(0) colloids in dichloromethane. (B) Iridium precursor used to synthesize Ir(0) nanoparticles in ionic liquid, trihexyltetradecylphosphium methylsulfonate [THTdP][MS]. (C) TEM image of the Ir nanoparticles. Small (d ~ 3.5 nm) and uniform nanoparticles were synthesized by the reduction of the Ir precursor in ethylene glycol under a hydrogen atmosphere. (D) Proposed catalytic cycle for the borylation of benzene. The ligands (L) in (D) represent [THTdP][MS] and [THTdP][MS] stabilized Ir(0) nanoparticles is considered the true catalyst (Zhu, et al., 2008). | 100 |
| Figure 4.12 | Bimetallic Au-Pd nanoparticle assembly for the cyclootrimerization of substituted acetylenes to highly substituted benzenes (Huang, et al., 2004)..... | 101 |
| Figure 4.13 | <i>In-situ</i> enzymatic saccharification of cellulose in a water-ionic liquid mixture. The IL disrupts the hydrogen-bonding network to increase the (Kamiya, et al., 2008)..... | 104 |
| Figure 4.14 | Metal foams can be synthesized via decomposition of bi(tetrazolato)amines like a) Fe BTA. b) Fully 3-D foams can be formed via this method as shown. (Tappan, et al., 2010) | 108 |

| | | |
|-------------|---|-----|
| Figure 4.15 | Self-assembly of polymetallic cluster nodes (left; top: $m_4\text{-oxo } M_4O(-CO_2)_6$; bottom: $M_2(-CO_2)_4$ paddlewheel) and organic linkers (right) yielding metal–organic frameworks (center) (Farrusseng <i>et al.</i> , 2009). | 110 |
| Figure 4.16 | Metal Organic Framework materials may operate as catalysts in two ways: a) via the generation of unsaturated metal connecting points as active catalytic sites or b) via the use of functional groups in the bridging ligands as active catalysts (Ma, <i>et al.</i> , 2010) | 112 |
| Figure 4.17 | Scheme for multicomponent photocatalyst for hydrogen production. (Youngblood, <i>et al.</i> , 2009)..... | 113 |
| Figure 4.18 | The mechanism of photocaging a catalyst via a photochemically induced cleavage of an inhibitor (or ligand) from its complex with the catalyst (Stoll, <i>et al.</i> , 2010)..... | 114 |
| Figure 4.19 | Photoswitchable azobenzene-based piperidines as light-controlled general base catalysts a) Concept of light-triggered reversible steric shielding of a basic/nucleophilic site. b) Chemical structures of investigated azobenzene and stilbene derivatives with optimized substitution pattern to achieve high on/off ratios (Stoll, <i>et al.</i> , 2010). | 116 |

TABLES

| | | |
|-----------|---|----|
| Table 2.1 | Translation of the 12 Green Chemistry Principles to Green Nanosynthesis (from Dahl <i>et al.</i> , 2007) | 6 |
| Table 2.2 | Pros and Cons of Surfactant-directed and Solvent-controlled Nonaqueous Liquid-Phase Routes to Metal Oxide Nanoparticles (from Pinna and Niederberger, 2008)..... | 7 |
| Table 2.3 | Examples of Metal Oxide Nanoparticles Synthesized by Surfactant-free Nonaqueous Sol–Gel Routes (from Pinna and Niederberger, 2008). | 8 |
| Table 2.4 | Alkylation of Acetophenone with Benzyl Alcohol at 150°C After 20 h in the Presence of Various Metal Nitride and Carbide Nanoparticles (from Yao <i>et al.</i> , 2009)..... | 10 |
| Table 2.5 | Space–time–yield for MOF synthesis (from Czaja <i>et al.</i> , 2009) | 26 |
| Table 2.6 | Examples of Reactions Catalyzed by MOFs (from Czaja <i>et al.</i> , 2009)..... | 27 |
| Table 4.1 | Parameter for the synthesis of Pt nanoparticles from di-block co-polymer micelles of poly(styrene)-block-poly(2-vinylpyridine) in toluene. The diameter of the nanoparticles was measured after supporting them on anatase TiO_2 and calcination for 2.5 h at 773 K (Croy, <i>et al.</i> , 2007)..... | 93 |

| | | |
|-----------|---|----|
| Table 4.2 | Reaction Orders, with Respect to Hydrogen Pressure for Different Palladium/Ionic Liquid/Activated Carbon Cloth (Pd/IL/ACC) Catalysts (Virtanen, et al., 2007). | 99 |
| Table 4.3 | [2+2+2] alkyne cyclotrimerization reactions catalyzed by RS-Au-PdCl ₂ nanoparticles in [bmim][PF ₆] under microwave irradiation (Lin, et al., 2008). 102 | |

SCHEMES

| | | |
|------------|--|-----|
| Scheme 4.1 | Hydroformylation of 1-octene to the corresponding linear and branched aldehyde over a xanthene-based diphosphine modified organometallic Rh catalyst..... | 103 |
| Scheme 4.2 | Tetrahydropyranylation of alcohols using 3,4-dihydro-2 <i>H</i> -pyran to the corresponding tetrahydropyranyl ethers in the presence of acids such as <i>p</i> -toluenesulfonic acid (TsOH), pyridinium <i>p</i> -toluenesulfonate (PPTS) and triphenylphosphine hydrobromide (TPP.HBr) using 3,4-dihydro-2 <i>H</i> -pyran in [bmim][BF ₄] or [bmim][PF ₆]. | 105 |
| Scheme 4.3 | Condensation of veratryl alcohol to the supramolecular host compound, cyclotrimeratrylene was synthesized in the presence of H ₃ PO ₄ in tributylhexylammonium bis(trifluoromethylsulfonyl)amide | 105 |
| Scheme 4.4 | Asymmetric Diels-Alder reaction between the dienophile (acryloyl oxazolidinone) and cyclopentadiene in the IL, 1,3-dibutylimidazolium tetrafluoroborate in the presence of a chiral catalyst, copper bisoxazoline based chiral Lewis acid. | 106 |

The Catalytic Advances Program (CAP)

CAP is the industry's leading membership-driven program which serves as an information resource for R&D and commercial organizations in the catalyst and process industries. Companies that join CAP combine their resources to jointly explore the world's most promising catalytic technologies. Members receive three in-depth **CAP Technical Reports** which are written by leading scientists and experienced industry professionals in areas selected by the membership (via ballot); weekly *CAP Communications* (delivered via e-mail), which provide the latest updates on technical breakthroughs, commercial events and exclusive development opportunities; and attendance at the CAP Annual Meeting.

The *Catalytic Advances Program (CAP)* is available on a membership basis from The Catalyst Group Resources (TCGR). For further details, please contact John J. Murphy at John.J.Murphy@catalystgrp.com or +1.215.628.4447.



P.O. Box 680
Spring House, PA 19477 U.S.A
ph: +1.215.628.4447
fax: +1.215.628.2267
website: www.catalystgrp.com