

ADVANCES IN CATALYSIS WITH PRECIOUS METALS

A technical investigation
commissioned by the members of the
Catalytic Advances Program (CAP)

Client Private
September 2013



The Catalytic Advances Program (CAP)

The Catalytic Advances Program (CAP) is an information resource for research and development organizations in the petroleum, chemical, and polymer industries. By the direction of the member companies (through balloting and other interactive means), the program delivers a range of timely and insightful information and analyses which are accessible exclusively to members and protected by confidentiality agreements. The objective is to provide a technical update on commercially viable advances in catalysis as well as benchmark commercial advances in catalysis and process technology.

Members receive three in-depth **CAP Technical Reports** which are written and peer reviewed by leading scientists and experienced industry professionals in areas selected by the membership (via ballot); weekly *CAP Communications* (delivered via e-mail) which provide the latest updates on technical breakthroughs, commercial events and exclusive development opportunities; and attendance at the **CAP Annual Meeting**.

The **Catalytic Advances Program (CAP)** is available on a membership basis from The Catalyst Group Resources (TCGR). For further details, please contact Matthew A. Colquitt at Matthew.A.Colquitt@catalystgrp.com or +1.215.628.4447 (x1130).



P.O. Box 680
Spring House, PA 19477 U.S.A
ph: +1.215.628.4447
fax: +1.215.628.2267
website: www.catalystgrp.com

CONTENTS

EXECUTIVE SUMMARY	xvii
1 INTRODUCTION	1
1.1 HISTORICAL BACKGROUND.....	1
1.2 COST PERFORMANCE OF PRECIOUS METALS.....	3
1.3 PRECIOUS METALS RECYCLING.....	5
1.4 AUTHORS & CONTRIBUTORS.....	7
1.5 REFERENCES.....	8
2. ADVANCES IN CATALYSIS WITH PLATINUM GROUP METALS	11
2.1 THE USE OF PLATINUM GROUP METALS IN CATALYTIC CONVERTORS.....	11
2.2 PLATINUM GROUP METALS CATALYSTS FOR THE PREFERENTIAL OXIDATION OF CO IN THE PRESENCE OF H ₂	13
2.3 ADVANCES IN CATALYSIS BY PGMS.....	15
2.3.1 Catalysis by Platinum.....	15
2.3.1.1 Catalytic wet air oxidation.....	15
2.3.1.2 Asymmetric hydrogenation.....	16
2.3.1.3 Water gas shift.....	17
2.3.1.4 Steam reforming biomass.....	18
2.3.2 Catalysis by Palladium.....	18
2.3.2.1 Oxidation reactions.....	18
2.3.2.2 Biomass conversion.....	23
2.3.2.3 Isomerization reactions.....	26
2.3.2.4 Hydrogenation reactions.....	28
2.3.2.5 Other palladium catalyzed reactions.....	29
2.3.2.6 Promoters for palladium.....	33
2.3.3 Catalysis by Ruthenium.....	34
2.3.3.1 Hydrogenation of CO.....	34
2.3.3.2 Other hydrogenation reactions.....	37
2.3.3.3 Oxidation reactions.....	38
2.3.3.4 Ammonia synthesis.....	41

2.3.3.5	Biomass conversion.....	42
2.3.4	Catalysis by Iridium.....	44
2.3.4.1	Selective hydrogenation.....	45
2.3.4.2	Selective ring opening (SRO).....	46
2.3.5	Catalysis by Osmium.....	47
2.3.5.1	Olefin oxidation.....	48
2.3.6	Catalysis by Rhodium.....	48
2.3.6.1	Steam reforming.....	48
2.3.6.2	CO oxidation.....	50
2.3.6.3	Pentanediol from biorenewables.....	51
2.4	TECHNICAL HURDLES.....	52
2.5	CONCLUSIONS AND RECOMMENDATIONS.....	53
2.6	REFERENCES.....	54
3.	ADVANCES IN CATALYSIS WITH GOLD AND SILVER.....	63
3.1	CATALYSIS BY GOLD.....	63
3.1.1	Selective Oxidation and Oxygen/ Metal Interaction.....	63
3.1.1.1	Alcohol oxidation.....	63
3.1.1.2	CO oxidation.....	64
3.1.2	Biomass Conversion.....	65
3.1.3	Water Gas Shift Reaction: $\text{CO} + \text{H}_2\text{O} \rightarrow \text{CO}_2 + \text{H}_2$	66
3.1.4	Effect of Gold Concentration.....	68
3.1.5	Gold as a Promoter.....	69
3.1.6	Unsupported Gold as a Catalyst.....	69
3.1.7	C-H Activation.....	71
3.2	CATALYSIS BY SILVER.....	72
3.2.1	Oxidation Reactions.....	73
3.2.1.1	Epoxidation.....	73
3.2.1.2	Alcohol oxidation/dehydrogenation.....	77
3.2.1.3	CO oxidation in the presence and absence of H_2	80
3.2.1.4	Hydrocarbon oxidation.....	82
3.2.1.5	Oxidative coupling of methane.....	83
3.2.2	Automotive Exhaust Catalysts.....	83
3.2.2.1	Selective catalytic reduction of NO_x	84
3.2.2.2	Soot oxidation.....	86

3.2.3	Hydrogenation Reactions	88
3.2.4	Reactions of Nitrogen Containing Compounds.....	89
3.3	BIMETALLIC CATALYSTS.....	89
3.3.1	AuPd.....	89
3.3.2	AuPt.....	101
3.3.3	AuCu	104
3.4	TECHNICAL HURDLES.....	108
3.5	CONCLUSIONS AND RECOMMENDATIONS.....	109
3.6	REFERENCES.....	111
4.	INDEX.....	123

FIGURES

Figure 1.1	Number of years remaining of rare and precious metal reserves if consumption and disposal continues at present rate.....	4
Figure 2.1	Continuous catalytic wet air oxidation of an ammonium acetate effluent as a function of time on stream in a trickle-bed reactor. Initial total nitrogen (TN) concentration (●), TN concentration (◆) over 3%Pt/ZrO ₂ (contact time = 1.055 h g _{Pt} g _N ⁻¹), and 3%Pt/TiO ₂ (▲, □) (contact time = 1.055 h g _{Pt} g _N ⁻¹ and 2.11 h g _{Pt} g _N ⁻¹ , respectively).....	16
Figure 2.2	Selective oxidation of benzyl alcohol over Pd/Al ₂ O ₃ and Pd/TiO ₂ at 393K	21
Figure 2.3	a) Effect of metal composition on cyclooctene conversion. Reaction conditions: catalyst (0.12 g), cis-cyclooctene (10 mL, 0.077 mol), TBHP (0.01mL, 1.03 x 10 ⁻⁴ mol), temperature 80 1C, glass reactor, 24 h, atmospheric pressure. b) Effect of metal composition on the selectivity of cyclooctene oxidation. Reaction conditions: catalyst (0.12 g), cis-cyclooctene (10 mL, 0.077 mol), TBHP (0.01mL, 1.03 x 10 ⁻⁴ mol), temperature 80 1C, glass reactor, 24 h, 10 ⁻⁴ mol) temperature 80 °C, glass reactor, 24h, atmospheric pressure. Epoxide (◆), alcohol (■), ketone (▲), hydroperoxide (●).....	22
Figure 2.4	Conversion and product distribution from furfural over 0.5%Pd/SiO ₂ as a function of W/F. Temp 230°C, H ₂ /feed ratio 25, H ₂ pressure 1atm, TOS 15min	24
Figure 2.5	Conversion and product distribution from 2-methyl pentanal over 5%Pd/SiO ₂ catalyst. Temp 125 °C, H ₂ /feed ratio 12, H ₂ pressure – 1 atm, TOS 15min	25

Figure 2.6	Effect of metal composition on the product yield of crotyl alcohol conversion. Reaction conditions: Catalyst (0.12 g), crotyl alcohol (10 mL, 0.12 mol), temperature 80 °C, TBHP (0.01 mL, 1.03×10^{-4} mol), 24 h, atmospheric pressure. Butyraldehyde (☐), crotonaldehyde (■), 3-buten-1-ol (□), acetic acid (▣), lactone epimer 1 (⊠), 4-hydroxy-2-butanone (▧), lactone epimer 2 (⊞), butyric acid (⊞).	28
Figure 2.7	Scanning transmission electron micrographs of (a) Pd/Al ₂ O ₃ and (b) Pd/La–Al ₂ O ₃ ; (c) particle size distribution of Pd in the catalysts, where atomic average particle size is the size of a Pd particle with the same dispersion as the sample; (d) high-resolution STEM of Pd/La–Al ₂ O ₃ showing lattice images of the Pd nanoparticles and atomically dispersed species on the alumina, which are La and Pd single atoms.	31
Figure 2.8	Catalyst performance for diesel oxidation catalysts as reported by BASF	32
Figure 2.9	Plots of (A) activity and (B) selectivity as a function of reaction temperature for CO preferential oxidation in the presence of hydrogen on PdPt alloy nanocages of 8, 12, and 20 nm in size, Pd nanocubes of 18 nm in size, and the commercial Pt/C catalyst.	34
Figure 2.10	The DFT derived activation energy for H-assisted CO activation on (111) terraces of Ru clusters at 1.55 monolayer CO* coverages is 165 kJ mol ⁻¹ . These values are much smaller than those for the unassisted CO dissociation (322 kJ mol ⁻¹).....	36
Figure 2.11	HMF has been identified as a versatile platform chemical obtained from biomass-derived carbohydrate which can be converted into a range of valuable chemicals.....	44
Figure 2.12	Aberration-corrected HAADF-STEM image of zeolite NaY-supported sample containing 10 wt % iridium after decarbonylation of the clusters: image showing Ir ₆ clusters with atomic resolution. Superimposed on the image next to two clusters are models of the cluster structures indicating the metal frameworks, which, because of electron-beam damage, are no longer octahedral, as they were in Ir ₆ (CO) ₁₆	45
Figure 2.13	Indan conversion (a) and single cleavage selectivity (b) vs. time-on-stream over different γ-Al ₂ O ₃ -supported nanoparticles at 336 °C and 0.1 MPa: Pd nanoparticles (●), Ir nanoparticles (◆), Ir nanoparticles (half amount) (◇), Pd ₂ Ir ₁ (△), Pd ₁ Ir ₁ (▲), Pd(c)Ir(s) (□), and industrial Pt–Ir catalyst (×).	46
Figure 2.14	Stages of the reaction between an Os-cluster and SWNT imaged by AC-HRTEM at 80 keV (a) with (b) the corresponding structural diagrams (numbers of metal atoms in diagrams are reduced for the sake of clarity). EDX spectroscopy confirms the identity of the metal clusters within the nanotubes as Os (c); Cu-peaks are due to the TEM specimen grid. High-magnification images showing details of key steps: π-bonding between an Os-cluster and the interior of the SWNT (d and e), σ-bonding between an Os-cluster and a defect on the SWNT (f), and π-bonding between an Os-cluster and the exterior of the SWNT (g).	47

Figure 2.15	Conversion (%) of ethanol over Rh/Al ₂ O ₃ at 773 K, 823 K and 873 K.....	50
Figure 3.1	Reaction scheme for biodiesel production	65
Figure 3.2	Temperature dependence of the CO conversion over the studied catalysts: (x) AuCe; (o) AuCeFeMA; (v) AuCeFeCP; (Δ) AuCeMnMA; (▲) AuCeMnCP; (μ) AuCeSnMA; (λ) AuCeSnCP.....	67
Figure 3.3	Rate per total mole of Au calculated at 120°C, 6.8% CO, 21.9% H ₂ O, 8.5% CO ₂ , and 37.4% H ₂ for Au/Al ₂ O ₃ (●) and Au/TiO ₂ catalysts (■) versus number average Au particle size. The vertical dimensions of the points represent the 15% error associated with rate measurement.....	68
Figure 3.4	Scanning electron microscopy (SEM) images of npAu samples. (a) <i>f</i> -NPG dealloyed at room temperature (rt) for 128 min in nitric acid under free corrosion conditions; (b) <i>a</i> -NPG dealloyed at rt for 15 min in nitric acid with an anodic potential of 1.0 V. Scale bars: 100 nm.	70
Figure 3.5	Particle size dependency of supported Au clusters for the oxidation of cyclohexane	72
Figure 3.6	The chlorohydrin process to make epoxides. A propylene chloronium complex is formed in the first step (1), followed by reaction with water to produce two propylene chlorohydrin isomers (2). The propylene chlorohydrin is then dehydrochlorinated using a base (usually calcium hydroxide) and the propylene oxide is steam-stripped from this reactor, to prevent base-catalyzed hydrolysis of the product.	73
Figure 3.7	The hydroperoxide process to make propylene oxide-styrene monomer. Ethylbenzene is oxidized to ethylbenzene hydroperoxide, which reacts with propylene to produce propylene oxide and α-phenyl ethanol. The α-phenyl ethanol is the dehydrated to styrene.	74
Figure 3.8	The oxametallacycle-surface interaction can be decomposed into C- and O–surface terms, which can be approximated by the interaction of methyl (CH ₃) and methoxy (CH ₃ O) radicals with the surface, respectively.....	75
Figure 3.9	Schematic reaction mechanism for propene epoxidation on silver. The adsorption of propene can result in hydrogen extraction to form an allylic intermediate or the propene oxametallacycle as branched (OMMP-B) or linear (OMMP-L) isomers.....	76
Figure 3.10	The mechanism of the ethanol to BD transformation is extremely complicated and is still subject of debate. However, one generally accepts the involvement of the following five principal steps: (1) acetaldehyde formation from ethanol; (2) aldol condensation of acetaldehyde to acetaldol; (3) dehydration of acetaldol to crotonaldehyde; (4) Meerwein–Ponndorf–Verley reaction between crotonaldehyde and ethanol to obtain crotyl alcohol and acetaldehyde; and (5) dehydration of crotyl alcohol to BD.	79

Figure 3.11	A scheme of the possible reaction pathways for CO selective oxidation over a silver catalyst pre-treated with oxygen at high temperature. The possible roles of oxygen in the feed gas are also demonstrated. There are competing CO ₂ pathways via surface oxygen (O _{weak}) and the more favorable pathway via sub-surface oxygen (O _γ). H ₂ cannot adsorb onto the surface and only reacts with surface oxygen at high temperature.	81
Figure 3.12	The structure of CeO ₂ -Ag catalysts with a rice ball morphology.....	87
Figure 3.13	Comparison of activity and reusability data for the direct synthesis of hydrogen peroxide for supported 0.5% Au/0.5%Pd/TiO ₂ catalysts prepared by sol-immobilization SIm (Cat-1), conventional impregnation CIm (Cat-2), and modified impregnation MIm methodologies (Cat-36). The modified impregnation catalysts tested are standard MIm dried only (Cat-3), reduced in 5% H ₂ /Ar at 400C/4 h (Cat-4) [hydrogen conversion and selectivity for this catalyst are 18% and 70% respectively], MIm prepared by NaCl excess (Cat-5), and MIm prepared by 2 M HCl (Cat-6). Reaction conditions: 5% H ₂ /CO ₂ and 25% O ₂ /CO ₂ , 50% H ₂ /O ₂ at 3.7 MPa, MeOH (5.6 g), H ₂ O (2.9 g), catalyst (0.01 g), 2C, 1200 rpm, 30 min.	96
Figure 3.14	Effect of the amount of “excess” HCl added during the synthesis of the 0.5%Au-0.5%Pd/TiO ₂ (MIm) catalyst on the resultant activity for the direct synthesis of hydrogen peroxide. In addition, the effect of the gas phase reduction temperature employed during the synthesis of a standard 0.5%Au-0.5%Pd/TiO ₂ (MIm) catalyst (using 0.58 M HCl) on the resultant activity for the direct synthesis of hydrogen peroxide is also plotted. Reaction conditions: 5% H ₂ /CO ₂ and 25% O ₂ /CO ₂ , 50% H ₂ /O ₂ at 3.7 MPa, MeOH (5.6 g), H ₂ O (2.9 g), catalyst (0.01 g), 2°C, 1200 rpm, 30 min. Key: (b) H ₂ O ₂ productivity vs HCl concentration; (9) H ₂ O ₂ productivity vs reduction temperature.....	97
Figure 3.15	(a) HRTEM image of a nanoflower consisting of a Au core and Pd petals. (b-d) Higher magnification images illustrate the polycrystalline nature of the Pd shell. White lines indicate the direction of lattice fringes in different crystalline domains.	98
Figure 3.16	Representative STEM HAADF images and corresponding XEDS analysis of particles from 2.5 wt% Au + 2.5 wt% Pd/TiO ₂ catalyst prepared by the physical mixing of acetate salts. The top row shows data from a typical large Au-rich particle (>50 nm in size), whereas the lower row shows data from a 2 nm AuPd nm alloy particle.....	99
Figure 3.17	The conversion of benzyl alcohol by catalysts prepared using various preparation techniques ▲ Au/TiO ₂ (PG), ■ AuPd/TiO ₂ (IMP), ○ AuPd/TiO ₂ (DP) , □ AuPd/TiO ₂ (PG), ◇ Pd/TiO ₂ (PG).....	99

Figure 3.18	Palladium shell thickness (nm) versus time (min) for the growth of faceted icosahedral Au–Pd core–shell nanocrystals. Time-resolved samples were obtained by quenching the reaction at 30, 60, 90, 120, and 180 min. The size distributions of each sample were measured and the shell thickness was calculated by subtracting the size of the gold seed. Standard size deviation is shown as error bars. Representative HRTEM images of each sample are shown in a)–f). Scale bars: 5 nm.	100
Figure 3.19	Catalytic properties of icosahedral Au–Pd core–shell nano-crystals with varying palladium shell thicknesses for the solvent-free oxidation of benzyl alcohol. a) Turnover frequency (TOF) versus palladium shell thickness.	101
Figure 3.20	Oxidation of 1,2-propanediol with air using the 0.5%Au + 0.5%Pt/KB-B catalyst. Reaction conditions: water (100 ml); 1,2-propanediol (0.6 M); NaOH/1,2-propanediol molar ratio = 1; 1,2-propanediol/total metal ratio = 5000; reaction time = 30–120 min; stirrer speed = 1000 rpm. Key: (■) 1,2-propanediol conversion; (◆) lactate selectivity	103
Figure 3.21	FDCA yield on different studied catalysts. Reaction conditions: 60 °C, 240 min, 10 bar of oxygen pressure 1 mmol of HMF, HMF : metal loading : NaOH molar ratio 1 : 0.01 : 4. Legend: □ monometallic Au on TiO ₂ , ■ bimetallic Au–Cu on TiO ₂	105
Figure 3.22	Effect of calcination or reduction on the stability of Au ₁ Cu ₁ /CeO ₂ catalyst under PROX reaction atmosphere. Inserts: Time on stream over 48h.	107
Figure 3.23	Amount of acetone formed during the aerobic oxidation of 2-propanol in the dark (black bars) or under visible-light irradiation (white bars; λ>450 nm; light intensity at 450–800 nm, 16.8 mWcm ⁻²).	108

TABLES

Table 1.1	Spot Precious Metal Price in US Dollars per Troy Ounce.....	5
Table 2.1	Catalytic performance of 0.5wt% Pd-containing catalysts supported on MnCeO _x	19
Table 2.2	Benzyl alcohol oxidation over Pd catalysts on different carbon supports	20
Table 2.3	Oxidation of HMF over supported Pt, Pd and Au catalysts.....	26
Table 2.4	The influence of O ₂ pressure on reaction and selectivity of HMF oxidation over Au/TiO ₂ catalyst	26
Table 2.5	Effect of preparation methods on (e)-2-hexen-1-ol oxidation	27
Table 2.6	Effect of support on (e)-2-hexen-1-ol oxidation.....	27

Table 2.7	SS-Ru catalyzed aerobic oxidation of alcohols to carbonyls.	39
Table 3.1	High activity of Au-Pd/nanotube catalysts for methane oxidation	72
Table 3.2	Comparison of catalytic activity for the oxidation of toluene in the absence of solvent with O ₂ . Catalyst mass varied between 0.2 and 0.6 g to give a substrate/metal molar ratio of 6500. Toluene, 10 to 20ml; stirring rate, 1500 rpm; TON calculated on the basis of the total metal. All metal catalysts were prepared using the sol immobilization method (18).	91
Table 3.3	Liquid-phase oxidation of toluene with TBHP as an oxidant at 80°C.	93
Table 3.4	Catalyst screening for oxidative imination of toluene	94
Table 3.5	Comparative catalytic activity of various Au-based catalysts for the liquid-phase oxidation of methane, with hydrogen peroxide either 1) added as co-reactant or 2) generated <i>in situ</i> from a H ₂ /O ₂ mixture under mild conditions.	95
Table 3.6	H ₂ O ₂ productivity over different supports: all catalysts are 2.5 wt% Au + 2.5 wt% Pd	100
Table 3.7	The effect of base on the oxidation of 1,2-propanediol using 1 wt% AuPt/carbon.....	102
Table 3.8	The oxidation of 1,2-propanediol using carbon supported nanoparticles in the absence of base at 40°C	102
Table 3.9	The oxidation of glycerol under base free conditions using selected gold bimetallic catalysts at low temperatures.	104
Table 3.10	Oxidation of various alcohol substrates over Au ₁ Cu ₁ /SiO ₂ catalyst.	106

SCHEMES

Scheme 2.1	Reactions involved in the DS process (i) synthesis, (ii) combustion, (iii) hydrogenation, (iv) decomposition	23
Scheme 3.1	The various oxidation pathways for toluene.	92